

Theory & Observations of Beam-beam Interactions in the Tevatron

Tanaji Sen

Fermi National Accelerator Laboratory


Tevatron Design Parameters

	Tevatron Run IIa
Luminosity [$\times 10^{31}$]	8.6
Bunch intensity [$\times 10^{11}$]	2.7/0.3
Normalized transverse emittance (p/\bar{p}) [95%, π mm-mrad]	20/15
RMS bunch length at top energy [cm]	37
RMS energy spread at top energy [$\times 10^{-4}$]	0.9
β^* [cm]	35
Beam-beam tune shift/IP [p/\bar{p}]	0.0014/0.01
Number of bunches	36
Total number of parasitics	72

Beam Losses in the Tevatron


	03/02	10/02	01/03	03/03
Protons/bunch at low-beta	140e9	170e9	180e9	205e9
Anti-protons/bunch at low-beta	7.5e9	22e9	20e9	23e9
P-loss at 150 GeV	23%	14%	16%	10%
Anti-proton-loss at 150 GeV	20%	9%	4%	4%
P-loss on ramp	7%	6%	9%	5%
Anti-proton-loss on ramp	14%	8%	12%	11%
Anti-proton-loss in squeeze	25%	5%	3%	2%

Dynamic aperture of protons at 150 GeV




Beam Separations at Injection

Injection: Helix ca. July 2002



Beta functions at all 138 parasitics: Injection energy



Beam-beam parameters vs Helix Angle

For round beams and large separations ($d \gg 1$), small amplitude parameters

$$\Delta\nu_x(0,0) \propto \frac{\cos 2\theta}{d^2}$$

$$\Delta\nu'_x(0,0) \propto \frac{\cos \theta(2 \cos 2\theta - 1)}{d^3} \eta_x$$

$$\Delta\nu_{min}(0,0) \propto \frac{\sin 2\theta}{d^2}$$


\Rightarrow

$\Delta\nu = 0$ along the diagonal

$\Delta\nu' = 0$ along 30° or the vertical axis.



$\Delta\nu_{min} = 0$ along the horizontal or vertical axis.

For arbitrary aspect ratio \Rightarrow



The tune shift is negative in the dark region and
vanishes along the border.



Small amplitude tune shifts & Tune footprints: Injection



Bunch to bunch tune spread $\Delta\nu_x \sim 0.005$.

Changes are small at the end of a train: A9-A12


Small amplitude chromaticities and coupling: Injection



Beam-beam chromaticity → some bunches more susceptible to synchro-betatron resonances, instabilities.

Changes are small towards the end of a train.

Anti-proton lifetimes at Injection: Stores 2420, 2502



Linear to Square Root Lifetime

Diffusion Equation (1D)

$$\frac{\partial f}{\partial \tau} = \frac{\partial}{\partial Z} \left[Z \frac{\partial f}{\partial Z} \right]$$

$$\tau \equiv \left(\frac{d\langle J \rangle}{dt} / J_a \right) t, \quad Z \equiv J / J_a.$$

Solution

$$N(\tau) = 2 \sum_n \frac{c_n}{\lambda_n} J_1(\lambda_n) e^{-\lambda_n^2 \tau / 4}$$

$$c_n = \frac{1}{J_1(\lambda_n)^2} \int_0^1 f_0(Z) J_0(\lambda_n \sqrt{Z}) dZ$$

If transverse ($\epsilon_{\perp} = 20\pi$ mm-mrad)


$$\Rightarrow \dot{\epsilon} \approx 16\pi \text{ mm} - \text{mrad/hr}$$

Very large.



If longitudinal (4 eV-sec bucket)

$$\Rightarrow \dot{\epsilon}_s \approx 1/3 \text{ eV} - \text{sec/hr}$$

Reasonable




Resonances at Injection



	Resonances (p, q) at 2σ	
	Largest	Others
Lattice driven	(2, -2)	(3,-1), (1,-1), (1,-3), (3,1), (4, 0)
Lattice and beam-beam driven	(3, 4)	(1, 6), (4, 3), (5, 2), (2, 5), (0, 7)


Anti-protons only - Beam Study



In this study, τ was well anti-correlated with the vertical emittance.

In typical stores, $1 \leq \tau(\bar{p}) \leq 10$ hours.



Anti-protons only - Beam Study



Loss of anti-protons during the ramp was very small $\sim 2\%$.


In typical stores, anti-proton losses during the ramp are $\sim 10\%$.

Anti-proton dynamic aperture at 150 GeV




Anti-proton dynamic aperture at 150 GeV


Anti-proton Dynamic Aperture at 150 GeV



Dynamic aperture of anti-proton bunch 1 at 150 GeV




Expected and Measured Intensity Drop: Store 2426




- The expected drop in intensity was calculated from the final bunch area (2D and 3D). The bunches are assumed to completely fill their dynamic aperture. The expected 3D loss and measured loss agree to within 2%.
- The largest difference in 2D and 3D areas was for bunch 1. This bunch had the greatest reduction in longitudinal emittance.
- This store occurred before the vertical dampers were restored. Since then, we have not seen this significant emittance shaving at injection.

Anti-proton lifetime at 150 GeV




Anti-proton lifetime vs Vert. Emittance at 150 GeV

Store 2420 (April 11, 2003)



Store 2502 (May 2, 2003) - Injection



Lifetime Simulations at 150 GeV

Parallel codes have been developed that run at NERSC

- A. Kabel(SLAC): Code **PlibB**


Fast evaluation of complex error function

- J.Qiang(LBNL): Code **Beambeam3D**

Uses a shifted Green's function approach


Both codes include the long-range interactions and transverse noise.

At small apertures, both predict lifetimes close to the observed lifetimes.





This predicts a qualitative increase in lifetime when the vertical chromaticity is dropped below 4 units.

Anti-proton lifetime at Injection - Several Stores




Dependence of anti-proton lifetime on proton intensities is low so far.

Anti-proton ramp efficiency in stores




Beam Separations at 980 GeV



Radial Separations at all beam-beam interactions - Collision helix



Beta functions at all beam-beam interactions: Collision





Small amplitude tune shifts & Tune footprints: Collision




Bunch 1 has lower ν_x , bunch 12 has lower ν_y .

Spread between bunches 2-11: $\Delta\nu_x \sim 0.0015$, $\Delta\nu_y \sim 0.001$.


Small amplitude chromaticities and coupling: Collision




Resonances at Collision




	Resonances (p, q) at 2σ	
	Largest	Others
Lattice driven	(2, -1)	(0, 3), (2, 1), (1, -3), (1,-2), (1, 2)
Lattice and beam-beam driven	(3, 4)	(3, 2), (1, 4), (2, 3), (5, 2), (2, 3)



Anti-proton emittance growth at Collision




Start of Store 2441.



Start of Store 2445: $\Delta\nu_y = -0.001$.

Anti-proton emittance growth at Collision



Vertical emittance growth rates in the two stores.

Losses vs Helix Size




Figure 1: Losses (at end of store) vs helix size.

Losses were observed at the end of a store by changing the helix size in both planes. Increasing the helix size to 10%

$$\tau(p) = 86\text{hrs} \rightarrow 68\text{hrs}$$

$$\tau(\bar{p}) = 43\text{hrs} \rightarrow 33\text{hrs}$$

Decreasing the helix size to 80% of original

$$\tau(p) \rightarrow 141\text{hrs}$$

$$\tau(\bar{p}) \rightarrow 67\text{hrs}$$

- Emittances did not change much at the end \Rightarrow tails were lost
- Tunes decreased < 0.002 down to 65% of original helix.

Sharp losses at the end were likely due to (7th, 12th) order resonances.

Status of Beam-beam effects

- **Injection**

- Limit anti-proton lifetimes to under 10 hrs
- No significant influence on protons

- **Ramp**

- Cause about 10% anti-proton losses
Anti-proton emittance growth during the ramp may be beam-beam related.
- Not much influence on protons

- **Squeeze**

- Anti-proton losses are low
- Proton losses are occasionally very high - causing quenches.

- **Collision**

- Anti-proton and proton lifetimes not much affected by beam-beam at present intensities in good stores.
- Occasionally have large emittance growth of anti-protons at start of store.
- Proton losses (thought to beam-beam related) can sometimes be higher than acceptable

Improvements

- Increasing the beam separations at all stages.
- Improving the alignment in the Tevatron.
- Smaller beam emittances.
- Operating with lower chromaticities (together with octupoles).
- Improved IR optics, e.g. local decoupling.
- Different bunch patterns.
- Active compensation of beam-beam effects

Acknowledgements

BNL: W. Fischer, F. Pilat, V. Ptitsyn

CERN: J.P. Koutchouk, F. Schmidt, F. Zimmermann

FNAL: Y. Alexahin, J. Annala, D. Capista, M. Church, D. Edwards,
B. Erdelyi, N. Gelfand, B. Hanna, I. Kourbanis, J. Johnstone,
V. Lebedev, P. Lebrun, M. Martens, R. Moore, S. Pordes, V. Shiltsev,
M. Syphers, D. Still, C.Y. Tan, M. Xiao, P. Yoon, X.L. Zhang

LBNL: J. Qiang

SLAC: A. Kabel